FIG.1.

Viral titres vs amount of DNA used for transfection




FIG.2






FIG.3



3/6


FIG.4

	Envelope and genome-containing particle Envelope containing empty particle		© © ©	\text{\tint{\text{\tin}\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\text{\tex{\tex	OO OO OO	9 90 0 00 0 00 0 00	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	HO A A A	<pre></pre>
	© ©								
Legend	• Envelope Receptor Genome-containing core	Bald genome-containing particle Bald empty particle	(A) (A) (A) (B) (B) (B) (B) (B) (B) (B) (B) (B) (B	(A)	1.100000	000000	0 0 0 0 0 0 0 + \$ag/gag-tot	1.100000	<pre>0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0</pre>



5°GGAUCCCGAUCUUCUGAUGAGUCCGUGAGGACGAAACGAGUUCCAUG CAGGUGCUGAUGAGUCCGUGAGGACGAAACCUCUGCGCCCUGCUCUGAU GAGUCCGUGAGGACGAAACGUGCCUGAAUUC-3°

SUBSTITUTE SHEET (RULE 26)



EGS Based on Tyrosyl t-RNA Precursor

6/6

FIG.6B